

DEGENERATE FLAG VARIETIES OF TYPE A AND C ARE SCHUBERT VARIETIES

GIOVANNI CERULLI IRELLI, MARTINA LANINI

ABSTRACT. We show that any type A or C degenerate flag variety is in fact isomorphic to a Schubert variety in an appropriate partial flag manifold.

1. INTRODUCTION AND MAIN RESULT

Appeared for the first time in the 19th Century to encode questions in enumerative geometry, flag varieties and their Schubert varieties had been intensively studied since then, constituting an important investigation object in topology, geometry, representation theory and algebraic combinatorics. In the years, several variations of these varieties have been considered (affine flag and Schubert varieties, Kashiwara flag varieties, matrix Schubert varieties, toric degenerations of flags, ...). Among them, we want to focus on a class introduced recently by E. Feigin in [7]: the degenerate flag varieties. These are flat degenerations of (partial) flag manifolds and turned out to be very interesting from a representation theoretic and geometric point of view. For instance, they can be used to determine a q -character formula for characters of irreducible modules in type A [9, 11] and C [12, 13]. As for the geometry, degenerate flag varieties share several properties with Schubert varieties: they are irreducible, normal locally complete intersections with terminal and rational singularities [7, 9, 13]. In this work we show that any degenerate flag variety of type A or C not only has a lot in common with Schubert varieties, but it is actually isomorphic to a Schubert in an appropriate partial flag variety. In short:

Theorem 1.1. *Degenerate flag varieties of type A and C are Schubert varieties.*

This result is based on the realization of degenerate flag varieties in terms of linear algebra, which is due to E. Feigin in type A [8, Theorem 2.5] and to E. Feigin, M. Finkelberg and P. Littelmann in type C [13, Theorem 1.1]. This description does not use any further information on the geometry of such varieties, and hence the theorem provides an independent proof of their geometric properties such as normality, irreducibility, rational singularities, cellular decomposition, which have been established in [7], [8], [9] and [13] by direct analysis.

We now state the precise version of Theorem 1.1 in the case of complete flags of type A (in Section 3 we discuss the case of partial flags, while in Section 4 we discuss the symplectic case). Let $n \geq 1$ and $B \subset SL_{2n}$ be the subgroup of upper triangular matrices. For a weight λ of SL_{2n} , let P_λ be its stabilizer. Let $\omega_1, \dots, \omega_{2n}$ be the fundamental weights and let $P := P_{\omega_1 + \omega_3 + \dots + \omega_{2n-1}}$ from now on. The Weyl group of SL_{2n} is Sym_{2n} (the symmetric group on $2n$ letters) and P corresponds to the subgroup W_J of Sym_{2n} generated by the traspositions $J = \{(2i, 2i+1)_{i=1, \dots, n-1}\}$. The variety SL_{2n}/P is naturally identified with the set of partial flags $W_1 \subset W_2 \subset \dots \subset W_n$ in \mathbb{C}^{2n} such that $\dim(W_i) = 2i - 1$.

The subgroup B acts on SL_{2n}/P (by left multiplication) and its orbits give the Bruhat decomposition:

$$(1.1) \quad SL_{2n}/P = \coprod_{\tau \in \text{Sym}_{2n}^J} B\tau P/P,$$

where Sym_{2n}^J is the set of permutations τ in Sym_{2n} such that $\tau(2i) < \tau(2i+1)$, for $i = 1, \dots, n-1$. This is the set of minimal length representatives for the cosets in Sym_{2n}/W_J . For a permutation $\tau \in \text{Sym}_{2n}^J$, let \mathcal{C}_τ be the corresponding Schubert cell in SL_{2n}/P , that is $B\tau P/P$, and denote by $X_\tau = \overline{B\tau P/P}$ its closure, that is the associated Schubert variety. Then each Schubert cell \mathcal{C}_τ has exactly one point which is fixed by the action of the subgroup of diagonal matrices $T \subseteq B$, namely

$$\langle e_{\tau(1)} \rangle < \langle e_{\tau(1)}, e_{\tau(2)}, e_{\tau(3)} \rangle < \dots < \langle e_{\tau(1)}, e_{\tau(2)}, e_{\tau(3)}, \dots, e_{\tau(2n-1)} \rangle.$$

(For a collection of vectors \mathbf{v} of a complex vector space, we always denote by $\langle \mathbf{v} \rangle$ the subspace spanned by \mathbf{v} .) Let $\sigma = \sigma_n \in \text{Sym}_{2n}$ be the permutation defined as

$$(1.2) \quad \sigma_n(r) = \begin{cases} k & \text{if } r = 2k, \\ n+1+k & \text{if } r = 2k+1. \end{cases}$$

For example, for $n = 5$ the permutation σ is given by

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 \\ 6 & 1 & 7 & 2 & 8 & 3 & 9 & 4 & 10 & 5 \end{pmatrix}.$$

Notice that $\sigma \in \text{Sym}_{2n}^J$, indeed $\sigma(2i) = i < \sigma(2i+1) = n+1+i$ for $1 \leq i \leq n-1$.

Let $\mathcal{F}l_{n+1}^a$ denote the complete degenerate flag variety associated with SL_{n+1} (see Section 2 for a definition of such a variety). In [3] it is shown that $\mathcal{F}l_{n+1}^a$ is acted upon by the maximal torus T of SL_{2n} (this is recalled in Section 2).

We are now ready to state the precise version of Theorem 1.1 in the case of complete flags (the general result for partial flags is Theorem 3.1).

Theorem 1.2. *There exists a T -equivariant isomorphism of projective varieties*

$$\zeta : \quad \mathcal{F}l_{n+1}^a \xrightarrow{\simeq} X_\sigma \subset SL_{2n}/P$$

where σ is the permutation given in (1.2) and $P = P_{\omega_1 + \omega_3 + \dots + \omega_{2n-1}}$.

We notice that since the isomorphism ζ is T -equivariant, it is possible to compute the stalks of the local T -equivariant intersection cohomology of $\mathcal{F}l_{n+1}^a$ by using the parabolic analogue of Kazhdan-Lusztig polynomials, defined by Deodhar in [6]. This answers a question posed in [3] (and it was the original motivation for this project). Another corollary of the theorem is that the median Genocchi number $h_n = \chi(\mathcal{F}l_{n+1}^a)$ (see [8]) has another interpretation: it is the number of elements $\tau \in \text{Sym}_{2n}^J$ which are smaller than σ in the (induced) Bruhat order.

As the anonymous referee has kindly pointed out, it would be interesting to analyze Theorem 1.1 in the light of the recent theory of Favourable Modules [10]. Moreover, She/He also asks what is the interplay between the Borel subgroup $B \subset \tilde{G}$ and the “degenerate group” G^a (see [8]), where $\tilde{G} = SL_{2n}$ (resp. $\tilde{G} = Sp_{4n-2}$) and $G = SL_{n+1}$ (resp. $G = Sp_{2n}$). Indeed both G and \tilde{G} act on the corresponding degenerate flag varieties (see Theorem 1.2 and 4.1 for the action of \tilde{G} and [8, 13] for the action of G^a). This will be the subject of a future project.

The paper is organized as follows: in Section 2 we prove Theorem 1.2, in Section 3 we discuss its analogue for partial degenerate flags and in Section 4 we prove the analogous results for type C. In the Appendix we prove that the desingularization of the degenerate flag varieties provided in [9] coincides with a Bott-Samelson resolution of the Schubert variety X_σ .

2. PROOF OF THEOREM 1.2

Given an integer $n \geq 1$, let $\mathcal{F}l_{n+1}^a$ denote the complete degenerate flag variety associated with SL_{n+1} . In [8, Theorem 2.5] it is proven that $\mathcal{F}l_{n+1}^a$ can be realized as follows: let $\{f_1, \dots, f_{n+1}\}$ be an ordered basis of a complex vector space $V \simeq \mathbb{C}^{n+1}$ and let $\text{pr}_k : V \rightarrow V$ be the linear projection along the line spanned by f_k , i.e. $\text{pr}_k(\sum a_i f_i) = \sum_{i \neq k} a_i f_i$. Then there is an isomorphism

$$\mathcal{F}l_{n+1}^a \simeq \{(V_1, \dots, V_n) \in \prod_{i=1}^n \text{Gr}_i(V) \mid \text{pr}_{i+1}(V_i) \subset V_{i+1} \forall i = 1, \dots, n-1\}.$$

For convenience of notation, up to an obvious change of basis of V , we prefer to realize $\mathcal{F}l_{n+1}^a$ as follows:

$$(2.1) \quad \mathcal{F}l_{n+1}^a \simeq \{(V_1, \dots, V_n) \in \prod_{i=1}^n \text{Gr}_i(V) \mid \text{pr}_i(V_i) \subset V_{i+1} \forall i = 1, \dots, n-1\}.$$

Let $\{e_1, \dots, e_{2n}\}$ be an ordered basis of a vector space $W \simeq \mathbb{C}^{2n}$. For any $i = 1, 2, \dots, n$, we consider the coordinate subspace $U_{n+i} := \langle e_1, \dots, e_{n+i} \rangle \subseteq W$ and the surjection $\pi_i : U_{n+i} \twoheadrightarrow V$ defined on the basis vectors as

$$(2.2) \quad \pi_i(e_k) = \begin{cases} 0 & \text{if } 1 \leq k \leq i-1, \\ f_k & \text{if } i \leq k \leq n+1, \\ f_{k-n-1} & \text{if } n+2 \leq k \leq n+i. \end{cases}$$

and extended by linearity to U_{n+i} . This induces a chain of embeddings of projective varieties

$$\begin{aligned} \text{Gr}_i(V) &\hookrightarrow \text{Gr}_{2i-1}(U_{n+i}) \hookrightarrow \text{Gr}_{2i-1}(W) \\ U &\longmapsto \pi_i^{-1}(U) \longmapsto \pi_i^{-1}(U) \end{aligned}$$

We call $\zeta_i : \text{Gr}_i(V) \hookrightarrow \text{Gr}_{2i-1}(W)$ the concatenation of the above maps. We hence have a diagonal embedding

$$(2.3) \quad \begin{aligned} \zeta : \prod_{i=1}^n \text{Gr}_i(V) &\longrightarrow \prod_{i=1}^n \text{Gr}_{2i-1}(W) \\ (V_1, V_2, \dots, V_n) &\longmapsto (\zeta_1(V_1), \zeta_2(V_2), \dots, \zeta_n(V_n)) \end{aligned}$$

Let us show that ζ restricts to a map $\mathcal{F}l_{n+1}^a \rightarrow SL_{2n}/P$. We consider a point $(V_1, \dots, V_n) \in \mathcal{F}l_{n+1}^a \subset \prod_{i=1}^n \text{Gr}_i(V)$; thus, $\text{pr}_i(V_i) \subset V_{i+1}$ for any $i = 1, \dots, n-1$. We notice that π_{i+1} coincides with $\text{pr}_i \circ \pi_i$ on $U_{n+i} \subset U_{n+i+1}$. Denoting by $W_i := \zeta_i(V_i)$, we get

$$W_i \subseteq \pi_{i+1}^{-1} \pi_{i+1}(W_i) = \pi_{i+1}^{-1} \text{pr}_i \pi_i(W_i) = \pi_{i+1}^{-1} \text{pr}_i(V_i) \subseteq \pi_{i+1}^{-1}(V_{i+1}) = W_{i+1}.$$

Therefore ζ restricts to an embedding $\zeta : \mathcal{F}l_{n+1}^a \hookrightarrow SL_{2n}/P$.

We now recall the action of the maximal torus $T \subset SL_{2n}$ on $\mathcal{F}l_{n+1}^a$ defined in [2, Section 3.1]. Let T_0 be a maximal torus of $GL_{n+1}(\mathbb{C})$. Up to a change of basis, we assume that T_0 acts on V by rescaling the basis vectors f_i 's. This induces a diagonal action of n copies $T_0^{(1)} \times \dots \times T_0^{(n)}$ of T_0 on the direct sum $V^{(1)} \oplus \dots \oplus V^{(n)}$ of n copies of V . More precisely we endow every copy $V^{(i)}$ with a basis $\{f_1^{(i)}, \dots, f_{n+1}^{(i)}\}$ and the torus acts by rescaling the $f_k^{(i)}$'s. We consider the linear map $\text{pr}_i : V^{(i)} \rightarrow V^{(i+1)}$ defined on the basis vectors by sending $f_k^{(i)}$ to $f_k^{(i+1)}$ for $k \neq i$, and $f_i^{(i)}$ to zero, and extended by linearity. We define $T_1 \subset \prod_{i=1}^n T_0^{(i)}$ to be the maximal subgroup such that each projection $\text{pr}_i : V^{(i)} \rightarrow V^{(i+1)}$ is T_1 -equivariant. It can be checked that

T_1 has dimension $2n$ and hence T_1 is isomorphic to a maximal torus of $GL_{2n}(\mathbb{C})$. More explicitly, an element $\underline{\lambda} = (\lambda_1, \dots, \lambda_{2n}) \in T_1$ acts by

$$(2.4) \quad \underline{\lambda} \cdot f_k^{(i)} := \begin{cases} \lambda_k f_k^{(i)} & \text{if } i \leq k \leq n+1 \\ \lambda_{n+1+k} f_k^{(i)} & \text{if } 1 \leq k \leq i-1 \end{cases}$$

Moreover, since the action of T_0 on V induces an action on each Grassmannian $\text{Gr}_i(V)$, then the action of $T_0^{(1)} \times \dots \times T_0^{(n)}$ on $V^{(1)} \oplus \dots \oplus V^{(n)}$ induces an action of T_1 on the product of Grassmannians $\prod_{i=1}^n \text{Gr}_i(V^{(i)}) = \prod_{i=1}^n \text{Gr}_i(V)$. Since each projection pr_i is T_1 -equivariant, this action descends to an action on $\mathcal{F}l_{n+1}^a$. Notice that the action of a point $\underline{\lambda} \in T$ on $\mathcal{F}l_{n+1}^a$ coincides with the action of any of its multiples; we hence see that $T := T_1 \cap SL_{2n}$ also acts on $\mathcal{F}l_{n+1}^a$.

We now prove that the map $\zeta : \mathcal{F}l_{n+1}^a \hookrightarrow SL_{2n}/P$ is T -equivariant. The maximal torus T in SL_{2n} acts on W (and hence on each Grassmannian $\text{Gr}_k(W)$) by rescaling the basis vectors e_k 's : given $\underline{\lambda} = (\lambda_1, \dots, \lambda_{2n}) \in T$

$$(2.5) \quad \underline{\lambda} e_k := \lambda_k e_k.$$

From (2.4) and (2.5) it follows that each map π_i is T -equivariant and hence each ζ_i is T -equivariant and hence ζ itself is T -equivariant.

We now describe the image $\zeta(\mathcal{F}l_{n+1}^a) \simeq \mathcal{F}l_{n+1}^a$. We claim that it is given by

$$(2.6) \quad Y_n := \left\{ W_1 \subset W_2 \subset \dots \subset W_n \mid \begin{array}{l} \bullet \dim W_i = 2i - 1 \\ \bullet \langle e_1, e_2, \dots, e_{i-1} \rangle \subset W_i \\ \bullet W_i \subset \langle e_1, \dots, e_{n+i} \rangle \end{array} \right\} \subset SL_{2n}/P.$$

Indeed, $\zeta(\mathcal{F}l_{n+1}^a)$ is clearly contained in Y_n ; viceversa, given a flag $W_\bullet := (W_1 \subset W_2 \subset \dots \subset W_n)$ in Y_n , then by definition $\ker \pi_i \subset W_i \subset U_{n+i}$ and hence $W_i = \pi_i^{-1}(\pi_i(W_i)) = \zeta_i(\pi_i(W_i))$. It follows that $W_\bullet = \zeta((\pi_1(W_1), \dots, \pi_n(W_n))) \in \text{Im } \zeta$. It remains to show that $(\pi_1(W_1), \dots, \pi_n(W_n)) \in \mathcal{F}l_{n+1}^a$. This is immediately verified as follows: $\text{pr}_i(\pi_i(W_i)) = \pi_{i+1}(W_i) \subseteq \pi_{i+1}(W_{i+1})$, for any $i = 1, \dots, n-1$.

In order to show that $Y_n \cong X_\sigma$, we observe that for any $i = 1, \dots, n$ we have

$$\#\{l \leq 2i-1 \mid \sigma(l) \leq k\} = \begin{cases} k & \text{if } 1 \leq k \leq i-1, \\ i-1 & \text{if } i-1 \leq k \leq n, \\ i-1+k-n & \text{if } n+1 \leq k \leq n+i, \\ 2i-1 & \text{if } n+i \leq k \leq 2n. \end{cases}$$

It follows that for a partial flag $W_\bullet \in SL_{2n}/P$, condition $\langle e_1, e_2, \dots, e_{i-1} \rangle \subseteq W_i \subseteq \langle e_1, e_2, \dots, e_{n+i} \rangle$ is equivalent to

$$(2.7) \quad \dim(W_i \cap \langle e_1, e_2, \dots, e_k \rangle) \geq \#\{l \leq 2i-1 \mid \sigma(l) \leq k\}$$

for any $i = 1, \dots, n$ and $k = 1, \dots, 2n$. By [14, Corollary of the proof of Proposition 7, §10.5], X_σ is precisely the locus of partial flags in SL_{2n}/P satisfying (2.7). This concludes the proof of Theorem 1.2.

Remark 2.1. Theorem 1.2 and its proof have a nice and clean interpretation in terms of quivers, in the spirit of [2], [3] and [4].

3. PARABOLIC CASE

In this section we discuss the parabolic analogue of Theorem 1.2. Recall the vector space $V \simeq \mathbb{C}^{n+1}$ with basis $\{f_1, \dots, f_{n+1}\}$ and let $\mathbf{d} = (d_i)_{i=1}^s$ be a collection of positive integers $1 \leq d_1 < d_2 < \dots < d_s \leq n$. For any pair of indices $1 \leq i < j \leq n$ we consider the linear map $\text{pr}_{i,j} : V \rightarrow V$ defined by $\text{pr}_{i,j} = \text{pr}_{j-1} \circ \dots \circ \text{pr}_{i+1} \circ \text{pr}_i$

where pr_i is the projection along f_i as before. Then, following [8, Theorem 2.5], the partial degenerate flag variety $\mathcal{F}l_{\mathbf{d}}^a$ is given by

$$\mathcal{F}l_{\mathbf{d}}^a \simeq \{(V_1, \dots, V_s) \in \prod_{l=1}^s \text{Gr}_{d_l}(V) \mid \text{pr}_{d_l, d_{l+1}}(V_l) \subset V_{l+1}\}.$$

The maximal torus $T \subset SL_{2n}$ acts on $\mathcal{F}l_{\mathbf{d}}^a$, in a similar way as for complete flags (see [2]). Let $\lambda := \omega_{2d_1-1} + \omega_{2d_2-1} + \dots + \omega_{2d_s-1}$ and let $P = P_{\lambda}$ be the corresponding parabolic subgroup in SL_{2n} . The variety SL_{2n}/P is naturally identified with the variety of partial flags $W_1 \subset \dots \subset W_s \subset W$ such that $\dim W_i = 2d_i - 1$ ($i = 1, 2, \dots, s$). We introduce the sets $K := \{1, 2, \dots, 2n\} \setminus \{2d_i - 1 \mid i = 1, 2, \dots, s\}$, $J := \{(k, k+1) \mid k \in K\}$, and the subgroup W_J of Sym_{2n} generated by J . We have the Bruhat decomposition

$$SL_{2n}/P \simeq \coprod_{\tau} B\tau P/P$$

where this time τ runs over the set of minimal length representatives for the cosets in Sym_{2n}/W_J . This set corresponds to the permutations $\tau \in \text{Sym}_{2n}$ such that $\tau(2d_i) < \tau(2d_i + 1) < \dots < \tau(2d_{i+1} - 1)$. We denote by $X_{\tau} = \overline{B\tau P/P}$ the corresponding Schubert variety. Let $\sigma_{\mathbf{d}}$ be the minimal length representative of the coset $\sigma_n W_J \in \text{Sym}_{2n}/W_J$ (σ_n is defined in (1.2)); explicitly, $\sigma_{\mathbf{d}}$ is given by

$$(3.1) \quad \sigma_{\mathbf{d}}(k) = \begin{cases} k - d_i & \text{if } k \in \{2d_i, \dots, d_i + d_{i+1} - 1\}, \\ n + 1 + k - d_{i+1} & \text{if } k \in \{d_i + d_{i+1}, \dots, 2d_{i+1} - 1\}, \end{cases}$$

with the conventions $d_0 := 0$ and $d_{s+1} := n + 1$. For example, for $n = 8$ and $\mathbf{d} = (2, 5, 7)$, we have

$$\sigma_{\mathbf{d}} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 & 14 & 15 & 16 \\ 1 & 9 & 10 & 2 & 3 & 4 & 11 & 12 & 13 & 5 & 6 & 14 & 15 & 7 & 8 & 16 \end{pmatrix}$$

Notice that for $\mathbf{d} = (1, 2, \dots, n)$, the permutations $\sigma_{\mathbf{d}}$ and σ_n (1.2) coincide.

Theorem 3.1. *There exists a T -equivariant isomorphism*

$$\zeta : \mathcal{F}l_{\mathbf{d}}^a \xrightarrow{\simeq} X_{\sigma_{\mathbf{d}}} \subset SL_{2n}/P_{\lambda}.$$

Proof. Recall the vector space $W \simeq \mathbb{C}^{2n}$ with basis $\{e_1, \dots, e_{2n}\}$ and the surjections $\pi_i : U_{n+i} \twoheadrightarrow V$ defined in (2.2) for $i = 1, 2, \dots, n$. The map ζ is defined by sending $(V_1, \dots, V_s) \in \mathcal{F}l_{\mathbf{d}}^a$ to the tuple $(W_1, \dots, W_s) \in SL_{2n}/P_{\lambda}$ given by $W_i := \pi_{d_i}^{-1}(V_i)$. It can be checked in the same way as in Section 2, that the image of ζ consists of partial flags $W_1 \subset W_2 \subset \dots \subset W_s$ such that $\dim W_i = 2d_i - 1$ and $\langle e_1, e_2, \dots, e_{d_i-1} \rangle \subseteq W_i \subseteq \langle e_1, \dots, e_{n+d_i} \rangle$. The proof now finishes as for Theorem 1.2. \square

4. SYMPLECTIC CASE

In this section we state and prove the analogue of Theorem 1.2 in the case of the symplectic group. In order to fix notation, we start with a brief overview about symplectic flag varieties (see e.g. [16, Chapter 6]). We consider a positive integer $n \geq 1$ and a complex vector space $W \simeq \mathbb{C}^{2n}$ of dimension $2n$ with ordered basis $\{e_1, e_2, \dots, e_{2n}\}$. We fix the bilinear form $b_W[\cdot, \cdot]$ on W given by the following $2n \times 2n$ matrix

$$(4.1) \quad E = \begin{pmatrix} 0 & J \\ -J & 0 \end{pmatrix}$$

where J is $n \times n$ anti-diagonal matrix with entries $(1, 1, \dots, 1)$, as usual. In particular the form is non-degenerate and skew-symmetric. Moreover $e_k^* = e_{2n+1-k}$, for $k = 1, \dots, 2n$. The group Sp_{2n} consists of those matrices A in SL_{2n} which leave

invariant the given form, i.e. $b_W[Av, Aw] = b_W[v, w]$ for every $v, w \in W$. More explicitly, we consider the involution $\iota : \mathrm{SL}_{2n} \rightarrow \mathrm{SL}_{2n}$ which sends a matrix A to the matrix $E({}^t A)^{-1} E^{-1}$; then the group Sp_{2n} consists of ι -invariant matrices. The advantage of choosing the form as above is that the intersection $B \cap \mathrm{Sp}_{2n} = B^\iota \subset \mathrm{SL}_{2n}$ consisting of ι -fixed upper triangular matrices, is indeed a Borel subgroup of Sp_{2n} whose maximal torus is precisely the subgroup $T^\iota = T \cap \mathrm{Sp}_{2n}$ of ι -invariant diagonal matrices.

The parabolic subgroup $P = P_{\omega_1 + \dots + \omega_{2n-1}}$ of SL_{2n} considered in Section 1 is mapped into itself by ι and the group of fixed points $Q := P^\iota = P \cap \mathrm{Sp}_{2n}$ is a parabolic subgroup of Sp_{2n} . The projective variety Sp_{2n}/Q can be described as follows: for a subspace $U \in \mathrm{Gr}_k(W)$ we denote by $U^\perp \in \mathrm{Gr}_{2n-k}(W)$ the orthogonal space of U in W . The map

$$(4.2) \quad \iota_k : \mathrm{Gr}_k(W) \rightarrow \mathrm{Gr}_{2n-k}(W) : U \mapsto U^\perp$$

is an isomorphism of projective varieties. The variety SL_{2n}/P sits inside the product $\prod_{i=1}^n \mathrm{Gr}_{2i-1}(W)$ and we consider the involution (still denoted by ι)

$$(4.3) \quad \iota := \prod_{i=1}^n \iota_{2i-1} : \prod_{i=1}^n \mathrm{Gr}_{2i-1}(W) \rightarrow \prod_{i=1}^n \mathrm{Gr}_{2i-1}(W)$$

The involution ι restricts to an involution on SL_{2n}/P and the variety $\mathrm{Sp}_{2n}/Q = (\mathrm{SL}_{2n}/P)^\iota$ consists of ι -invariant flags.

Moreover, the involution ι (on SL_{2n}) induces an involution on the symmetric group Sym_{2n} as follows: it sends $\tau \mapsto \iota(\tau)$, where $\iota(\tau)(r) := 2n+1 - \tau(2n+1-r)$, for $r = 1, \dots, 2n$. The Weyl group of Sp_{2n} coincides with the subgroup Sym_{2n}^ι of ι -fixed elements. The element $\sigma_n \in \mathrm{Sym}_{2n}$ defined in (1.2) is easily seen to be fixed by ι and it hence belongs to the Weyl group of Sp_{2n} . The left action of B^ι on Sp_{2n}/Q induces the Bruhat decomposition:

$$\mathrm{Sp}_{2n}/Q = \coprod_{\tau \in (\mathrm{Sym}_{2n}^\iota)^\iota} B^\iota \tau Q/Q.$$

Each Schubert cell $B^\iota \tau Q/Q$ coincides with the set of ι -fixed points \mathcal{C}_τ^ι of the Schubert cell \mathcal{C}_τ of SL_{2n} and the same holds for each Schubert variety, $Z_\tau = \overline{B^\iota \tau Q/Q} = X_\tau^\iota$ (cf. [16, Proposition 6.1.1.2]).

We now state the analogue of Theorem 1.2 in type C. We denote by SpFl_{2m}^a the complete degenerate flag variety associated with Sp_{2m} (see below for a definition).

Theorem 4.1. *There exists a T^ι -equivariant isomorphism of projective varieties*

$$(4.4) \quad \zeta : \mathrm{SpFl}_{2m}^a \xrightarrow{\cong} X_{\sigma_n}^\iota \subset \mathrm{Sp}_{2n}/Q$$

where $n := 2m - 1$, σ_n is the permutation given in (1.2) and $Q = P^\iota$ as above.

In Section 4.1 we prove Theorem 4.1 and in Section 4.2 we state and prove its parabolic analogue.

4.1. Proof of Theorem 4.1. Fix an integer $m \geq 1$, a complex vector space V of dimension $2m$ with basis $\{f_1, \dots, f_{2m}\}$ and a non-degenerate skew-symmetric bilinear form $b_V[\cdot, \cdot]$ on V such that

$$(4.5) \quad f_k^* = \begin{cases} f_{2m-1-k} & \text{if } 1 \leq k \leq 2m-2, \\ f_{2m} & \text{if } k = 2m-1, \end{cases}$$

so that $V = \langle f_1, \dots, f_{m-1}, f_{m-1}^*, \dots, f_1, f_{2m-1}, f_{2m-1}^* \rangle$. We define $n := 2m - 1$, so that V has dimension $n+1$ as in the previous sections. The degenerate flag variety \mathcal{Fl}_{n+1}^a sits inside the product of Grassmannians $\prod_{i=1}^n \mathrm{Gr}_i(V)$. It can be checked that the map $\iota = \prod_i \iota_i : \prod_{i=1}^n \mathrm{Gr}_i(V) \rightarrow \prod_{i=1}^n \mathrm{Gr}_i(V)$ (where ι_i is defined in (4.2))

restricts to a map from $\mathcal{F}l_{n+1}^a$ to itself, and the fixed points form the degenerate symplectic flag variety associated with Sp_{2m} [13, Proposition 4.7], i.e.

$$(4.6) \quad \mathrm{Sp}\mathcal{F}l_{2m}^a = (\mathcal{F}l_{n+1}^a)^\iota.$$

Thus Theorem 4.1 will follow once we show that the diagram

$$(4.7) \quad \begin{array}{ccc} \mathcal{F}l_{n+1}^a & \xrightarrow{\iota} & \mathcal{F}l_{n+1}^a \\ \zeta \downarrow & & \downarrow \zeta \\ X_{\sigma_n} & \xrightarrow{\iota} & X_{\sigma_n} \end{array}$$

commutes, where the vertical arrows denote the T -equivariant isomorphism provided by Theorem 1.2 and the horizontal arrow in the bottom is induced by the involution (4.3). In Section 2 we proved that the isomorphism ζ is the restriction of the map $\zeta : \prod_{i=1}^n \mathrm{Gr}_i(V) \rightarrow \prod_{i=1}^n \mathrm{Gr}_{2i-1}(W)$ given in (2.3). In order to prove (4.7), it is enough to show that the following diagram

$$(4.8) \quad \begin{array}{ccc} \prod_{i=1}^n \mathrm{Gr}_i(V) & \xrightarrow{\iota} & \prod_{i=1}^n \mathrm{Gr}_i(V) \\ \zeta \downarrow & & \downarrow \zeta \\ \prod_{i=1}^n \mathrm{Gr}_{2i-1}(W) & \xrightarrow{\iota} & \prod_{i=1}^n \mathrm{Gr}_{2i-1}(W) \end{array}$$

commutes. We therefore need to check that for every point $(V_i)_{i=1}^n \in \prod_{i=1}^n \mathrm{Gr}_i(V)$ and for every $i = 0, \dots, m-1$, we have

$$(4.9) \quad \zeta_{m-i}(V_{m-i})^\perp = \zeta_{m+i}(V_{m-i}^\perp).$$

Recall that for every $i = 1, \dots, n$, $\zeta_i(V_i) := \pi_i^{-1}(V_i)$, where $\pi_i : U_{n+i} \rightarrow V$ is the map given in (2.2) and U_{n+i} is the coordinate subspace of W generated by e_1, e_2, \dots, e_{n+i} . We prove the following (stronger) statement: for every $i = 0, \dots, m-1$, $v \in U_{n+m-i}$ and $w \in U_{n+m+i}$ we have

$$(4.10) \quad b_V[\pi_{m-i}(v), \pi_{m+i}(w)] = b_W[v, w].$$

It is easy to verify that (4.10) implies (4.9): Indeed $\dim \zeta_{m-i}(V_{m-i})^\perp = 2m + 2i - 1 = \dim \zeta_{m+i}(V_{m-i}^\perp)$ and (4.10) implies at once that $\zeta_{m+i}(V_{m-i}^\perp) \subseteq \zeta_{m-i}(V_{m-i})^\perp$. We will prove (4.10) by induction on $i \geq 0$. For $i = 0$ we need to show that $\pi_m : U_{n+m} \rightarrow V$ is a map of symplectic spaces, i.e. for every $v, w \in U_{n+m}$ we have $b_V[\pi_m(v), \pi_m(w)] = b_W[v, w]$. This follows easily from the definitions: Indeed, for a given $k = 1, \dots, n$, the coordinate vector subspace U_{n+k} of W is given by $U_{n+k} = \langle e_1, \dots, e_n, e_n^*, \dots, e_{n-k+1}^* \rangle$. In particular, U_{n+m} is generated by $e_1, \dots, e_m, \dots, e_n, e_n^*, \dots, e_m^*$ and π_m is defined on the symplectic basis as follows

$$\pi_m(e_k) = \begin{cases} 0 & \text{if } 1 \leq k \leq m-1, \\ f_{n-k}^* & \text{if } m \leq k \leq n-1, \\ f_n & \text{if } k = n, \end{cases} \quad \pi_m(e_k^*) = \begin{cases} f_{n-k} & \text{if } m \leq k \leq n-1, \\ f_n^* & \text{if } k = n. \end{cases}$$

We hence assume that (4.10) is true for $i \geq 0$ and we prove it for $i+1$. In view of (4.5), the map $\mathrm{pr}_{m-1+k} : V \rightarrow V$ ($1 \leq k \leq m-1$) is the projection along the line spanned by the basis vector f_{m-k}^* and we denote $\mathrm{pr}_{(m-k)^*} := \mathrm{pr}_{m-1+k}$. We notice that the adjoint map pr_i^* of $\mathrm{pr}_i : V \rightarrow V$ is pr_{i^*} , i.e.

$$(4.11) \quad b_V[\mathrm{pr}_i(v), v'] = b_V[v, \mathrm{pr}_{i^*}(v')]$$

for every $v, v' \in V$. We have already observed that $\pi_{i+1} : U_{n+i+1} \rightarrow V$ restricted to $U_{n+i} \subset U_{n+i+1}$ coincides with $\mathrm{pr}_i \circ \pi_i$ and, using the notation just introduced,

this means that the following diagram

(4.12)

$$\begin{array}{cccccccccccccccc}
 V & \xrightarrow{\text{pr}_1} & V & \xrightarrow{\text{pr}_2} & \dots & \xrightarrow{\text{pr}_{m-1}} & V & \xrightarrow{\text{pr}_{m-1}^*} & V & \xrightarrow{\dots} & V & \xrightarrow{\text{pr}_2^*} & V & \xrightarrow{\text{pr}_1^*} & V \\
 \pi_1 \uparrow & & \pi_2 \uparrow & & & & \pi_{m-1} \uparrow & & \pi_m \uparrow & & \pi_{m+1} \uparrow & & \pi_{n-1} \uparrow & & \pi_n \uparrow \\
 U_{n+1} & \rightarrow & U_{n+2} & \rightarrow & \dots & \rightarrow & U_{n+m-1} & \rightarrow & U_{n+m} & \rightarrow & U_{n+m+1} & \rightarrow & \dots & \rightarrow & U_{2n-1} & \rightarrow & U_{2n}
 \end{array}$$

commutes (the chain of horizontal arrows in the bottom row is given by the canonical embeddings $U_{n+i} \hookrightarrow U_{n+i+1}$).

We can now prove (4.10). We write a non-zero element $w \in U_{n+m+(i+1)}$ as $w = \mu e_{n-m-i}^* + w'$ for some $w' \in U_{n+m+i}$ and some $\mu \in \mathbb{C}$; given $v \in U_{n+m-(i+1)}$ we need to compute $b_V[\pi_{m-(i+1)}(v), \pi_{m+(i+1)}(w)]$. Let us first deal with the case when $w' = 0$, i.e. $w = \mu e_{n-m-i}^*$: we have

$$\begin{aligned}
 b_V[\pi_{m-(i+1)}(v), \pi_{m+(i+1)}(w)] &= \mu b_V[\pi_{m-(i+1)}(v), \pi_{m+(i+1)}(e_{n-m-i}^*)] \\
 &= \mu b_V[\pi_{m-(i+1)}(v), f_{m+i}] \\
 &= \mu b_V[\pi_{m-(i+1)}(v), f_{m-1-i}^*].
 \end{aligned}$$

By writing $v = \sum_k c_k e_k$ in the symplectic basis $\{e_k\}$, since $\pi_{m-i-1}(e_{n-m-i}) = f_{n-m-i} = f_{m-1-i}$, we get

$$(4.13) \quad b_V[\pi_{m-(i+1)}(v), \pi_{m+(i+1)}(w)] = \mu c_{n-m-i} = b_W[v, \mu e_{n-m-i}^*] = b_W[v, w].$$

We now consider the case when $w' \neq 0$. In view of (4.11), (4.12), (4.13) and the induction hypothesis we get:

$$\begin{aligned}
 &b_V[\pi_{m-(i+1)}(v), \pi_{m+(i+1)}(w)] \\
 &= b_W[v, \mu e_{n-m-i}^*] + b_V[\pi_{m-(i+1)}(v), \pi_{m+(i+1)}(w')] \\
 &= b_W[v, \mu e_{n-m-i}^*] + b_V[\pi_{m-(i+1)}(v), \text{pr}_{m-i-1}^* \circ \pi_{m+i}(w')] \\
 &= b_W[v, \mu e_{n-m-i}^*] + b_W[\text{pr}_{m-i-1} \circ \pi_{m-i-1}(v), \pi_{m+i}(w')] \\
 &= b_W[v, \mu e_{n-m-i}^*] + b_V[\pi_{m-i}(v), \pi_{m+i}(w')] \\
 &= b_W[v, \mu e_{n-m-i}^*] + b_W[v, w'] \\
 &= b_W[v, w]
 \end{aligned}$$

as desired.

4.2. Parabolic case. We conclude by discussing the parabolic version of Theorem 4.1, which is the type C analogue of Theorem 3.1. Let $m \geq 1$ be a positive integer as in Section 4.1, and let $\mathbf{d} = (d_i)_{i=1}^s$ be a collection of positive integers $1 \leq d_1 < d_2 < \dots < d_s \leq 2m$ preserved by the map $d_i \mapsto 2m - d_i$. The involution $\iota = \prod \iota_i : \prod_{i=1}^s \text{Gr}_{d_i}(V) \rightarrow \prod_{i=1}^s \text{Gr}_{d_i}(V)$ is hence well-defined and restricts to a map from $\mathcal{F}_{\mathbf{d}}^a$ to itself. The fixed points form the partial degenerate symplectic flag variety $\text{Sp}\mathcal{F}_{\mathbf{d}}^a$ [13, Proposition 4.9], i.e. $\text{Sp}\mathcal{F}_{\mathbf{d}}^a = (\mathcal{F}_{\mathbf{d}}^a)^\iota$.

Let λ and P_λ as in Section 3, so that $X_{\sigma_{\mathbf{d}}} \subset \text{SL}_{2m}/P_\lambda$. Let $Q := P_\lambda^\iota$ be the parabolic subgroup of Sp_{2m} . The projective variety Sp_{2m}/Q coincides with the ι -fixed points of SL_{2m}/P_λ , i.e. $\text{Sp}_{2m}/Q = (\text{SL}_{2m}/P_\lambda)^\iota$. Moreover, since the permutation $\sigma_{\mathbf{d}}$ is fixed by ι , the corresponding Schubert variety in Sp_{2m}/Q is the variety of ι -fixed points $X_{\sigma_{\mathbf{d}}}^\iota$ of $X_{\sigma_{\mathbf{d}}}$. From the commutativity of Diagram (4.8), together with Theorem 3.1, we obtain the following result.

Theorem 4.2. *There exists a T^ι -equivariant isomorphism of projective varieties*


$$\zeta : \quad \text{Sp}\mathcal{F}_{\mathbf{d}}^a \xrightarrow{\cong} X_{\sigma_{\mathbf{d}}}^\iota \subset \text{Sp}_{2n}/Q$$

where $\sigma_{\mathbf{d}}$ is the permutation given in (3.1).

APPENDIX: DESINGULARIZATIONS

In [9] a resolution of the singularities of a type A degenerate flag variety is constructed. In view of Theorem 1.2 it is natural to ask if such a desingularization coincides with a Bott-Samelson resolution of the corresponding Schubert variety. In this section, we show that this is indeed the case. In order to state and prove the result we need a combinatorial model that we discuss in Section 4.3. In Section 4.4 we recall the construction of Feigin and Finkelberg. In Section 4.5 we construct a Bott-Samelson resolution of X_{σ_n} . In Section 4.6 we prove that the two desingularizations coincide (see Theorem 4.8). In the whole section $n \geq 1$ is a fixed integer.

4.3. The quiver Γ_n . We denote by Γ_n the quiver (i.e. the oriented graph) with $N := \binom{n+1}{2}$ vertices $\alpha_{i,j}$ with $1 \leq i \leq j \leq n$ and an oriented edge $\alpha_{i,j} \rightarrow \alpha_{i+1,j}$ (for every $1 \leq i < j \leq n$) and an oriented edge $\alpha_{i,j} \rightarrow \alpha_{i,j+1}$ (for $1 \leq i \leq j < n$). The quiver Γ_n is the famous Auslander-Reiten quiver of the equioriented type A quiver algebra (see e.g. [1]). For future reference we embed the quiver Γ_n into the decorated quiver $\tilde{\Gamma}_n$ obtained from Γ_n by adding $2n+1$ extra vertices $\alpha_{0,1}, \alpha_{0,2}, \alpha_{0,3}, \dots, \alpha_{0,n}, \alpha_{0,n+1}, \alpha_{1,n+1}, \alpha_{2,n+2}, \dots, \alpha_{n,2n}$ and oriented edges $\alpha_{0,i} \rightarrow \alpha_{1,i}$ (for $1 \leq i \leq n$) and $\alpha_{i,n} \rightarrow \alpha_{i,n+1}$ (for $1 \leq i \leq n$). For example for $n = 4$, the following is $\tilde{\Gamma}_4$ with the extra vertices highlighted:



Given an index $\ell = 0, 1, \dots, 2n$, we define the ℓ -th column of $\tilde{\Gamma}_n$ as the set of all vertices $\alpha_{i,j}$ such that $i+j-1 = \ell$. For example, the vertex $\alpha_{i,i}$ lies on the column $2i-1$. Similarly, given an index $r = 1, 2, \dots, n+2$, we define the r -th row of $\tilde{\Gamma}_n$ as the set of vertices $\alpha_{i,j}$ of $\tilde{\Gamma}_n$ such that $j-i+1 = r$. Notice that $\Gamma_n \subset \tilde{\Gamma}_n$ has $2n-1$ columns and n rows; the row at the bottom consists of the vertices $\{\alpha_{i,i}\}_{i=1}^n$ and the top row consists of the single vertex $\alpha_{1,n}$. Following [9] we order the vertices of Γ_n as $\beta_1 < \beta_2 < \dots < \beta_N$ so that every row is ordered from left to right and every element of a row is smaller than every element in the rows below. More precisely: we put $\beta_1 := \alpha_{1,n}$, and for $\beta_k := \alpha_{i_k, j_k}$ we put

$$\beta_{k+1} := \begin{cases} \alpha_{i_k+1, j_k+1} & \text{if } j_k < n, \\ \alpha_{1, n-i_k} & \text{if } j_k = n. \end{cases}$$

To illustrate: $\beta_1 = \alpha_{1,n}$, $\beta_N = \alpha_{n,n}$ and $\beta_{N-k} = \alpha_{n-k, n-k}$ for $k = 0, 1, \dots, n-1$. We sometimes need to extend this ordering on the vertices of Γ_n to an ordering on the vertices of $\tilde{\Gamma}_n$. We do this in the following way: we define $\beta_0 := \alpha_{0,1}$ and

$$\beta_{-k} := \alpha_{0, k+1}, \quad \beta_{-n-k} := \alpha_{k, n+1}$$

for every $1 \leq k \leq n$. We notice that the vertex β_{-m} lies on the m -th column of $\tilde{\Gamma}_n$ (for $0 \leq m \leq 2n$). We always use the following notation: the vertex β_k is equal to α_{i_k, j_k} and it lies on the column $\ell_k = \ell_{\beta_k} := i_k + j_k - 1$. Moreover for a vertex β_k

of Γ_n we denote by β_k^- and β_k^+ the vertices of $\tilde{\Gamma}_n$ given by:

$$(4.15) \quad \beta_k^- := \alpha_{i_k-1, j_k} \quad \beta_k^+ := \alpha_{i_k, j_k+1}.$$

By construction, both β_k^- and β_k^+ are *strictly smaller* than β_k and $\ell_{\beta_k^\pm} = \ell_{\beta_k} \pm 1$. We will also need the following notation: for a vertex β_k of Γ_n and a column index $\ell \in [1, 2n-1]$ of Γ_n we define the index

$$(4.16) \quad (\beta_k : \ell) = \max\{\beta_s \mid -2n \leq s \leq k, \beta_s \text{ lies on the } \ell\text{-th column}\}.$$

The following equalities are a direct consequence of the definition:

$$(4.17) \quad (\beta_k : \ell_k) = \beta_k$$

$$(4.18) \quad (\beta_k : \ell_k \pm 1) = \beta_k^\pm$$

$$(4.19) \quad (\alpha_{1,j} : \ell) = \beta_{-\ell} \quad \forall 0 \leq \ell < j,$$

$$(4.20) \quad (\beta_N : 2k-1) = \beta_{N-(n-k)} \quad \forall 1 \leq k \leq n$$

$$(4.21) \quad (\beta_k : t) = (\beta_{k-1} : t) \quad \forall t \neq \ell_k.$$

The following technical result will be used later.

Lemma 4.3. *For $1 \leq k \leq N$ and $1 \leq t \leq 2n-2$, either $(\beta_k : t+1) = (\beta_k : t)^+$ or $(\beta_k : t+1)^- = (\beta_k : t)$.*

Proof. Let $\beta_s := (\beta_k : t)$ and $\beta_r := (\beta_k : t+1)$. Then, β_s and β_r lie in different columns and either $\beta_s < \beta_r$ or $\beta_r < \beta_s$. If $\beta_s < \beta_r$ then $\beta_r^- = \beta_s$ by maximality of β_s . If $\beta_r < \beta_s$ then $\beta_r = \beta_s^+$ by maximality of β_r . \square

4.4. The desingularization of Feigin and Finkelberg. In this section we recall the construction of a desingularization $R_n \dashrightarrow \mathcal{F}l_n^a$ of the complete degenerate flag variety $\mathcal{F}l_n^a$ due to E. Feigin and M. Finkelberg [9]. We fix a complex vector space V of dimension $n+1$ and ordered basis $\{f_1, \dots, f_{n+1}\}$. For every $1 \leq k \leq N$ we consider the coordinate subspace V_{β_k} of V given by

$$(4.22) \quad V_{\beta_k} := \text{Im}(\text{pr}_{j_k-1} \circ \dots \circ \text{pr}_{i_k}) = \langle f_1, \dots, f_{i_k-1}, f_{j_k}, \dots, f_{n+1} \rangle$$

with the convention that $f_0 = 0$ so that $V_{1,j} = \langle f_j, \dots, f_{n+1} \rangle$ (as before, the parenthesis $\langle \dots \rangle$ denotes the \mathbf{C} -linear span of the collection of vectors enclosed within them and $\text{pr}_k : V \rightarrow V$ is the projection along the line generated by f_k). Notice that the coordinate subspace V_{β_k} is the analogous of W_{i_k, j_k} in [9] taking into account our notation (2.1). We decorate the collection $(V_{\beta_k})_{k=1}^N$ with the subspaces

$$V_{\beta_0} = V_{\beta_{-k}} := \{0\} \quad V_{\beta_{-n-k}} := V$$

for every $k = 1, \dots, n$. In [9, Definition 2.1], the variety R_n is defined as the variety of collections $(Z_{\beta_k})_{k=1}^N$ such that

$$(4.23) \quad Z_{\beta_k} \in \text{Gr}_{i_k}(V_{\beta_k}),$$

$$(4.24) \quad Z_{\beta_k^-} \subset Z_{\beta_k},$$

$$(4.25) \quad \text{pr}_{j_k}(Z_{\beta_k}) \subset Z_{\beta_k^+},$$

with the convention that $Z_{0,k} = \{0\}$ and $Z_{k,n+1} = V$ for $1 \leq k \leq n$. The variety R_n is a tower of \mathbf{P}^1 -fibrations: for every $1 \leq s \leq N$ we consider the variety $R_n(s)$ of “truncated” collections $(Z_{\beta_k})_{k=1}^s$ satisfying conditions (4.23), (4.24), (4.25). Then there is a natural epimorphism

$$p_n(s) : R_n(s+1) \dashrightarrow R_n(s) : (Z_{\beta_i})_{1 \leq i \leq s+1} \mapsto (Z_{\beta_i})_{1 \leq i \leq s}.$$

whose fiber is \mathbb{P}^1 . In particular R_n is a smooth projective variety and the map

$$p_n : R_n \rightarrow \mathcal{F}l_{n+1}^a : (Z_{\beta_k})_{k=1}^N \mapsto (Z_{i,i})_{1 \leq i \leq n}$$

is a desingularization of $\mathcal{F}l_{n+1}^a$ ([9]).

4.5. A Bott-Samelson resolution of X_{σ_n} . In this Section we discuss a Bott-Samelson resolution of the Schubert variety X_{σ_n} (see (1.2)). We choose the following reduced expression for the permutation $\sigma_n \in \text{Sym}_{2n}$:

$$(4.26) \quad \sigma_n = s_n \circ (s_{n-1} \circ s_{n+1}) \circ (s_{n-2} \circ s_n \circ s_{n+2}) \circ \cdots \circ (s_1 \circ s_3 \circ \cdots \circ s_{2n-1})$$

(s_i denotes the simple transposition $(i, i+1)$). In particular we see that the length of σ_n is $N = \binom{n+1}{2}$ and the simple reflections appearing in (4.26) are in bijection with the vertices of Γ_n as follows

$$\sigma_n = \tau_{\beta_1} \circ \tau_{\beta_2} \circ \cdots \circ \tau_{\beta_N}.$$

To illustrate: $\tau_{\beta_1} = s_n$, $\tau_{\beta_2} = s_{n-1}$, $\tau_{\beta_N} = s_{2n-1}$. By construction, $\tau_{\beta_k} = s_{\ell_k}$.

We fix a complex vector space W of dimension $2n$ with ordered basis $\{e_1, \dots, e_{2n}\}$. Given two complete flags

$$U_\bullet = (U_1 \subset U_2 \subset \cdots \subset U_{2n-1} \subset W), \quad W_\bullet = (W_1 \subset W_2 \subset \cdots \subset W_{2n-1} \subset W)$$

in SL_{2n}/B and an integer $k = 1, 2, \dots, 2n-1$, the pair (U_\bullet, W_\bullet) is said to be in *relative position* k if $U_i = W_i$ for every $i \neq k$. Let F_\bullet be the standard flag of W , i.e. $F_i = \langle f_1, \dots, f_i \rangle$, for every $0 \leq i \leq 2n$ ($F_0 := \{0\}$). Notice that the subspace F_{n+i} was denoted by U_{n+i} before and we freely use the two notations. We consider the variety

$$(4.27) \quad BS_n := \{(U_\bullet^{\beta_k})_{k=0}^N \in (SL_{2n}/B)^{N+1} \mid \begin{array}{l} U_\bullet^{\beta_0} = F_\bullet \text{ and } (U_\bullet^{\beta_{k+1}}, U_\bullet^{\beta_k}) \text{ is} \\ \text{in relative position } \ell_{k+1} \end{array}\}.$$

This variety is the Bott-Samelson variety associated with the reduced expression (4.26) of σ_n . In particular, the following result is well-known [15], [5].

Proposition 4.4. *The variety BS_n is smooth. The map*

$$(4.28) \quad \rho_n : BS_n \longrightarrow X_{\sigma_n} : (U_\bullet^{\beta_k})_{k=0}^N \longmapsto (U_{2k-1}^{\beta_N})_{k=1}^n$$

is a desingularization.

We now give a description of BS_n which will be used later.

Definition 4.5. Let B_n be the projective variety of collections $(U_{\beta_k})_{k=-2n}^N$ (indexed by the vertices of $\tilde{\Gamma}_n$) such that, for every $1 \leq k \leq N$ and $0 \leq t \leq 2n$,

$$(4.29) \quad U_{\beta_{-t}} = F_t,$$

$$(4.30) \quad U_{\beta_k} \in \text{Gr}_{\ell_k}(W),$$

$$(4.31) \quad U_{\beta_k} \subset U_{n+i_k},$$

$$(4.32) \quad U_{\beta_k^-} \subset U_{\beta_k} \subset U_{\beta_k^+}.$$

Proposition 4.6. *The map $\theta_n : BS_n \rightarrow B_n$ which sends a point $U_\bullet^{\beta_\bullet} = (U_\bullet^{\beta_k})_{k=0}^N$ to the collection $\theta_n(U_\bullet^{\beta_\bullet}) = (U_{\beta_k})_{k=-2n}^N$ defined by*

$$U_{\beta_k} := \begin{cases} U_{\ell_k}^{\beta_k} & \text{if } k \geq 0 \\ U_k^{\beta_0} & \text{if } k < 0 \end{cases}$$

is an isomorphism of projective varieties. The inverse $\psi = \theta_n^{-1}$ sends a point $(U_{\beta_k})_{k=-2n}^N$ of B_n to the collection of complete flags $(U_\bullet^{\beta_k})_{k=0}^N$ defined by

$$(4.33) \quad U_t^{\beta_k} := U_{(\beta_k : t)}$$

for every $1 \leq t \leq 2n-1$.

Proof. Let us prove that the map θ_n is well-defined. This is based on the following technical result.

Lemma 4.7. *Let $(U_{\bullet}^{\beta_k})_{k=0}^N$ be a point of BS_n . Then,*

(i) *For every $1 \leq k \leq N$, and $1 \leq t \leq 2n-1$ we have*

$$(4.34) \quad U_t^{\beta_k} = U_t^{(\beta_k:t)}$$

(ii) *For $1 \leq k \leq N$*

$$(4.35) \quad F_{j_k-1} \subset U_{\ell_k}^{\beta_k} \subset F_{n+i_k}$$

Proof. By definition, $U_t^{\beta_k} = U_t^{\beta_{k-1}}$ for $t \neq \ell_k$ and, in particular, (4.34) holds for $k=1$. By induction on $k \geq 1$, using (4.21), we get the desired (4.34).

If $\beta_k = \alpha_{1,j_k}$ then $\ell_k = j_k$ and $(\beta_k : j_k - 1) = \beta_{-(j_k-1)}$ (see (4.19)); in view of (4.34) we have $U_{\ell_k-1}^{\beta_k} = U_{\ell_k-1}^{\beta_0} = F_{\ell_k-1}$ and hence $F_{j_k-1} = U_{\ell_k-1}^{\beta_k} \subset U_{\ell_k}^{\beta_k}$. Similarly, if $\beta_k = \alpha_{i,n}$, then $\ell_k = n+i-1$ and $(\beta_k : \ell_k + 1) = \beta_k^+ = \beta_{-(n+i)}$; in view of (4.34), $U_{\ell_k+1}^{\beta_k} = U_{\ell_k+1}^{\beta_0} = F_{n+i}$ and hence $U_{\ell_k}^{\beta_k} \subset U_{\ell_k+1}^{\beta_k} = F_{n+i}$. We hence assume that both β_k^- and β_k^+ are vertices of Γ_n . In this case we have

$$U_{\ell_k-1}^{\beta_k^-} = U_{\ell_k-1}^{\beta_k} \subset U_{\ell_k}^{\beta_k} \subset U_{\ell_k+1}^{\beta_k} = U_{\ell_k+1}^{\beta_k^+}.$$

By induction on the row index, (using (4.15)) the desired (4.35) follows. \square

Given $U_{\bullet}^{\beta_{\bullet}} \in BS_n$, in view of Lemma 4.7 we have $U_{\ell_k}^{\beta_k} \in \text{Gr}_{\ell_k}(U_{n+i_k})$; moreover

$$U_{\ell_{\beta_k^-}}^{\beta_k^-} = U_{\ell_k-1}^{\beta_k} \subset U_{\beta_k} \subset U_{\ell_k+1}^{\beta_k} = U_{\ell_{\beta_k^+}}^{\beta_k^+}$$

and hence $\theta_n(U_{\bullet}^{\beta_{\bullet}})$ satisfies all the four conditions (4.29)–(4.32) and the map θ_n is well-defined and it is clearly algebraic.

With the help of Lemma 4.3 and in view of (4.34), one easily checks that the map ψ is well-defined. It is clearly algebraic and the composition with θ_n is the identity. \square

4.6. Main Result. We are now ready to state and prove the main result of the this appendix.

Theorem 4.8. *There exists an isomorphism $\psi_n : R_n \longrightarrow BS_n$ of projective varieties such that the following diagram*

$$(4.36) \quad \begin{array}{ccc} R_n & \xrightarrow{\psi} & BS_n \\ p_n \downarrow & & \downarrow \rho_n \\ \mathcal{F}l_{n+1}^a & \xrightarrow{\zeta} & X_{\sigma_n} \end{array}$$

commutes (where the map ζ is the one in Theorem 1.2).

Proof. Recall that for every $1 \leq i \leq j \leq n$ we have a commutative diagram

$$\begin{array}{ccc} U_{n+i} & \xrightarrow{\subset} & U_{n+j} \\ \pi_i \downarrow & & \downarrow \pi_j \\ V & \xrightarrow{\text{pr}_{j-1} \circ \dots \circ \text{pr}_i} & V \end{array}$$

and it follows immediately from (4.22) that

$$(4.37) \quad V_{\beta_k} = \pi_{j_k}(U_{n+i_k}).$$

Since $\dim \text{Ker}(\pi_{j_k}) = j_k - 1$ and hence $\ell_k = i_k + \dim \text{Ker}(\pi_{j_k})$, we have an embedding of projective varieties:

$$\begin{aligned} \zeta_{\beta_k} : \text{Gr}_{i_k}(V_{\beta_k}) &\hookrightarrow \text{Gr}_{\ell_k}(U_{n+i_k}), \\ Z &\longmapsto \pi_{j_k}^{-1}(Z). \end{aligned}$$

This induces an embedding

$$\zeta = \zeta_n := \prod_{k=1}^N \zeta_{\beta_k} : \prod_{k=1}^N \text{Gr}_{i_k}(V_{\beta_k}) \hookrightarrow \prod_{k=1}^N \text{Gr}_{\ell_k}(U_{n+i_k}).$$

In view of (4.23), the variety R_n is contained in $\prod_{k=1}^N \text{Gr}_{i_k}(V_{\beta_k})$ and we claim that the image $\zeta(R_n)$ is canonically isomorphic to the variety B_n (see Definition 4.5). Indeed let $U = (U_{\beta_k})_{k=1}^N = \zeta((Z_{\beta_k})_{k=1}^N) \in \zeta(R_n)$, i.e. $U_{\beta_k} := \pi_{j_k}^{-1}(Z_{\beta_k})$. We decorate the collection $U = (U_{\beta_k})_{k=1}^N$ with the subspaces $U_{\beta_{-t}} := F_t$, for $0 \leq t \leq 2n$. The decorated collection $\tilde{U} = (U_{\beta_k})_{k=-2n}^N$ satisfies (4.29), (4.30), (4.31) and (4.32), i.e. $\tilde{U} \in B_n$. Indeed, (4.29) is clearly satisfied, (4.30) follows from (4.23) and (4.31) follows from (4.24). Since $Z_{\beta_k} \subset V_{\beta_k} = \pi_{j_k}(U_{n+i_k})$, there exists $U \subset U_{n+i_k}$ such that $Z_{\beta_k} = \pi_{j_k}(U)$; in view of (4.25), $\text{pr}_{j_k}(Z_{\beta_k}) \subset Z_{\beta_k}^+$ and we have

$$\pi_{j_k+1}^{-1}(Z_{\beta_k}^+) \supset \pi_{j_k+1}^{-1} \text{pr}_{j_k}(Z_k) = \pi_{j_k+1}^{-1} \pi_{j_k+1}(U) \supseteq U = \pi_{j_k}^{-1}(Z_k)$$

and hence \tilde{U} satisfies (4.32).

On the other hand, let $(U_{\beta_k})_{k=-2n}^N$ be a point of B_n . We define $Z_{\beta_k} := \pi_{j_k}(U_{\beta_k})$ ($k = 1, 2, \dots, N$) and we check that $(Z_{\beta_k})_{k=1}^N \in R_n$, i.e. it satisfies (4.23), (4.24) and (4.25). In view of (4.37), $Z_{\beta_k} \in V_{\beta_k}$. Since $U_{\beta_k}^- \subset U_{\beta_k}$ we have $\text{Ker} \pi_{j_k} = F_{j_k-1} \subseteq U_{\beta_k}$ and hence $\dim Z_{\beta_k} = \ell_k - (j_k - 1) = i_k$. It follows that $Z_{\beta_k} \in \text{Gr}_{i_k}(V_{\beta_k})$ and (4.23) is satisfied. Since $U_{\beta_k}^- \subset U_{\beta_k}$, (4.24) follows. To prove (4.25), we notice that

$$\text{pr}_{j_k}(Z_{\beta_k}) = \text{pr}_{j_k} \circ \pi_{j_k}(U_{\beta_k}) = \pi_{j_k+1}(U_{\beta_k}) \subseteq \pi_{j_k+1}(U_{\beta_k}^+) = Z_{\beta_k}^+$$

and the claim is proved.

In view of Proposition 4.6, the map $\psi_n := \theta_n^{-1} \circ \zeta_n : R_n \rightarrow \text{BS}_n$ is hence an isomorphism of projective varieties. It remains to check that the diagram (4.36) commutes. Using (4.20), we get

$$\rho_n \circ \psi_n((Z_{\beta_k})_{k=1}^N) = (\pi_{j_k}^{-1}(Z_{\alpha_{k,k}}))_{k=1}^n.$$

On the other hand, $p_n((Z_{\beta_k})_{k=1}^N) = (Z_{\alpha_{k,k}})_{k=1}^n$, and

$$\zeta \circ p_n((Z_{\beta_k})_{k=1}^N) = (\pi_{j_k}^{-1}(Z_{\alpha_{k,k}}))_{k=1}^n$$

as desired. \square

Acknowledgments. The main part of this project took place during M.L. stay at the Department of Mathematics “Guido Castelnuovo” of “Sapienza-Università di Roma”. We thank that institution for the perfect working conditions. The work of M.L. was financed by DFG SPP1388 and “Teoria delle rappresentazioni e applicazioni, Progetto di Ateneo 2012, Sapienza Università di Roma”. The work of G.C.I. was financed by the national FIRB grant RBFR12RA9W. We thank Rocco Chirivì, Francesco Esposito and Paolo Bravi for helpful discussions on a previous version of the present paper. We are especially grateful to Corrado De Concini, Michael Finkelberg and Evgeny Feigin for their useful comments and to Peter Littelmann and Oksana Yakimova for several discussions about possible applications of this work.

REFERENCES

1. I. Assem, D. Simson, A. Skowronski, *Elements of the representation theory of associative algebras. Vol. 1. Techniques of representation theory*. London Mathematical Society Student Texts, 65. Cambridge University Press, Cambridge, 2006.
2. G. Cerulli Irelli, E. Feigin, M. Reineke, *Quiver Grassmannians and degenerate flag varieties*, Algebra & Number Theory **6** (2012), no. 1, 165–194. arXiv: 1106.2399.
3. G. Cerulli Irelli, E. Feigin, M. Reineke, *Degenerate flag varieties: moment graphs and Schröder numbers*, J. Algebraic Combin. **38** (2013), no. 1. arXiv:1206.4178.
4. G. Cerulli Irelli, E. Feigin, M. Reineke, *Desingularization of quiver Grassmannians for Dynkin quivers*, Adv. Math. **245** (2013), 182–207. arXiv:1209.3960.
5. M. Demazure, *Désingularisation des variétés de Schubert généralisées*, Ann. Sci. École Norm. Sup. **7** (1974), 53–88.
6. V. Deodhar, *On some geometric aspects of Bruhat orderings II. The parabolic analogue of Kazhdan-Lusztig polynomials*, J. Algebra **111** (1987), no. 2, 483–506.
7. E. Feigin, \mathbb{G}_a^M *degeneration of flag varieties*, Selecta Mathematica, **18**, no. 3 (2012), 513–537. arXiv:1007.0646.
8. E. Feigin, *Degenerate flag varieties and the median Genocchi numbers*, Mathematical Research Letters, **18** (2011), no. 6, 1–16. arXiv: 1101.1898.
9. E. Feigin, M. Finkelberg, *Degenerate flag varieties of type A: Frobenius splitting and BW theorem*, Math. Z., **275** (2013), no. 1-2, 55–77. arXiv:1103.1491.
10. E. Feigin, G. Fourier, P. Littelmann, *Favourable modules: Filtrations, polytopes, Newton-Okounkov bodies and flat degenerations*. Preprint (2013). arXiv: 1306.1292.
11. E. Feigin, G. Fourier, P. Littelmann, *PBW filtration and bases for irreducible modules in type A_n* . Transf. Groups **16** (2011), no. 1, 71–89. arXiv:1002.7694.
12. E. Feigin, G. Fourier, P. Littelmann, *PBW filtration and bases for symplectic Lie algebras*, Int. Math. Res. Not. **24** (2011), 5760–5784. arXiv:1010.2321.
13. E. Feigin, M. Finkelberg, P. Littelmann, *Symplectic degenerate flag varieties*, To appear in Canad. J. Math. <http://cms.math.ca/10.4153/CJM-2013-038-6>. arXiv:1106.1399.
14. W. Fulton, *Young tableaux*, London Mathematical Society Student Texts, **35**. Cambridge University Press, (1997).
15. H. C. Hansen, *On cycles in flag manifolds*, Math. Scand. **33** (1973), 269–274.
16. V. Lakshmibai, K.N. Raghavan, *Standard monomial theory. Invariant theoretic approach.*, Encyclopaedia of Mathematical Sciences, 137. Invariant Theory and Algebraic Transformation Groups, 8. Springer-Verlag, Berlin, 2008.

GIOVANNI CERULLI IRELLI: DIPARTIMENTO DI MATEMATICA. SAPIENZA-UNIVERSITÀ DI ROMA. PIAZZALE ALDO MORO 5, 00185, ROME (ITALY)

E-mail address: cerulli.math@gmail.com

MARTINA LANINI: DEPARTMENT MATHEMATIK, UNIVERSITÄT ERLANGEN-NÜRNBERG, CAUER-STRASSE 11, 91058 ERLANGEN (GERMANY)

E-mail address: lanini@math.fau.de